arXiv:1302.0624v2 [hep-ph] 26 Mar 2013

The Neutrino Mass Hierarchy from Nuclear Reactor Experiments

Emilio Ciuffoli, Jarah Evslin and Xinmin Zhang
TPCSF, IHEP, CAS, YuQuanLu 19B, Beijing 100049, China

10 years from now reactor neutrino experiments will attempt to determine which neutrino mass eigenstate is the most massive. In this letter we present the results of more than four million detailed simulations of such experiments, studying the dependence of the probability of successfully determining the mass hierarchy upon the analysis method, the neutrino mass matrix parameters, reactor flux models and, in particular, combinations of baselines. We show that a recently reported spurious dependence of the data analysis upon the high energy tail of the reactor spectrum can be removed by using a weighted Fourier transform. We determine the optimal baselines and corresponding detector locations. For most values of the CP-violating, leptonic Dirac phase δ, a degeneracy prevents NOνA and T2K from determining either δ or the hierarchy. We determine the confidence with which a reactor experiment can determine the hierarchy, breaking the degeneracy.

Last year the Daya Bay [1, 2] and RENO [3] experiments demonstrated that the neutrino mass matrix mixing angle θ_{13} is nonzero and as much as an order of magnitude larger than had been suspected several years ago. With the discovery that θ_{13} is nonzero, at least three qualitative questions remain to be answered in the standard model plus three massive neutrinos. First, it is not known whether the second or third neutrino flavor is the most massive. This choice is known as the neutrino mass hierarchy. Second, it is not known whether the leptonic sector has a non-zero CP-violating Dirac phase δ. Third, the most precise determination of the mixing angle, θ_{23}, by the MINOS collaboration is

$$\sin^2(2\theta_{23}) = 0.96 \pm 0.04. \tag{1}$$

This equation has eight distinct solutions, known as the octants of θ_{23}. These solutions can be divided into two families, corresponding to θ_{23} modulo 90° greater than or less than 45°.

The large value of θ_{13} means that an appearance experiment, searching for ν_e's in a beam of accelerator ν_μ's, may be sensitive to CP violation in the leptonic sector. The trouble is that, at the baselines of the T2K experiment in Japan and the NOνA experiment in the US, the CP phase δ is degenerate with the mass hierarchy, the octant of θ_{23} and the precise value of θ_{13}. Following the analysis in Ref. [5], by combining data from the appearance of ν_e's in a ν_μ beam with the appearance of $\overline{\nu}_e$'s in a $\overline{\nu}_\mu$ beam in its second three year run, given sufficient funding NOνA may be able to separate the octant and θ_{13}, which determine the total $\nu_e + \overline{\nu}_e$ appearance, from the hierarchy and δ, which roughly determine the difference $\nu_e - \overline{\nu}_e$. However for most values of δ even the combination of T2K and NOνA will not be able to break the degeneracy between δ and the hierarchy, and so will not be able to determine either.

Fortunately the large value of θ_{13} also implies that 1-3 neutrino oscillations, those with amplitudes proportional to $\sin^2(2\theta_{13})$, are large enough to be observed in reactor neutrino experiments at medium baselines (40-60 km). These oscillations are almost periodic in the inverse energy E, but their small aperiodicity may be used to determine the neutrino mass hierarchy [6] thus breaking the degeneracy at T2K and NOνA and so allowing a determination of the CP-violating Dirac phase. More precisely, combined with a hierarchy determination, the NOνA appearance mode can only determine $\sin(\delta)$ because the transformation $\delta \to \pi - \delta$ is degenerate with a slight shift in θ_{23} or θ_{13}. However the effective mass difference determined by a reactor experiment, which will be discussed below, differs from the δ-dependent atmospheric effective mass [7] determined by the disappearance channel of experiments such as MINOS and NOνA. This difference may yield a 1σ determination of $\cos(\delta)$. Such reactor experiments are not only possible but indeed *will* be performed within the next decade [8, 9].

In this letter we present the main results of a series of simulations of medium baseline reactor experiments. We will determine the reliability of the determinations of the hierarchy in such experiments and will also find the optimal baselines for their detectors. In most of our simulations we use the value of ΔM^2_{21} from Ref. [10], $\sin^2(2\theta_{12})$ from Ref. [11], $|\Delta M^2_{32}|$ determined by combining ν and $\overline{\nu}$ mass differences from Ref. [4] and $\sin^2(2\theta_{13})$ from [2]

$$\Delta M^2_{21} = 7.59 \times 10^{-5}\ \mathrm{eV}^2,\ \ \sin^2(2\theta_{12}) = 0.857$$
$$|\Delta M^2_{32}| = 2.41 \times 10^{-3}\ \mathrm{eV}^2,\ \ \sin^2(2\theta_{13}) = 0.089.$$

Since we hold these parameters fixed, our neural network is trained with the same parameters which are used to test it, which artificially inflates the chance p of correctly determining the hierarchy. We have also systematically studied the effects of changing these parameters, as will be described below.

We consider reactor experiments in which $\overline{\nu}_e$'s are detected via inverse β decay by a detector with a 20 kton target mass consisting of 10% hydrogen. This is the mass of the proposed Daya Bay II detector, but it is four times the mass of the proposed detector for RENO 50, and so to interpret the results below for RENO 50 one needs to multiply all times by a factor of four. We assume a detector resolution of $3\%/\sqrt{E(\mathrm{MeV})}$. We ignore backgrounds and assume a perfect detector energy response. As a result of these approximations our results are overly optimistic.

We have performed 3 and 6 year simulations using several reactor flux models with various cutoffs using fluxes arising from over 100 combinations of baselines. For each combination, we have have simulated 5,000 experimental runs with each hierarchy. The data analysis methods described below are then used to attempt to determine the hierarchy. We report the percentage p of experiments for which the hierarchy is determined correctly.

The determination of the neutrino mass hierarchy from 1-3 oscillations in the $\overline{\nu}_e$ spectrum proceeds as follows. At energies E/MeV greater than L/12 km, deviations from periodicity in $1/E$ are too small to be measured and the wavenumber k determines [7]

$$\Delta M^2_{\mathrm{eff}} = \cos^2(\theta_{12})|\Delta M^2_{31}| + \sin^2(\theta_{12})|\Delta M^2_{32}|. \quad (2)$$

At low energies the deviation from periodicity is large and it determines various combinations of $|\Delta M^2_{31}|$ and $|\Delta M^2_{32}|$ given in Ref. [12], for example the energy of the 16th oscillation peak is proportional to $|\Delta M^2_{31}|$. To determine the mass hierarchy one needs to combine two distinct combinations of $|\Delta M^2_{31}|$ and $|\Delta M^2_{32}|$, thus one must combine the high and low energy parts of the spectrum.

The most studied algorithm which determines the hierarchy given a reactor $\overline{\nu}_e$ spectrum is that of Ref. [13]. One first finds the Fourier transform of the measured spectrum, as suggested in Ref. [14]. 1-3 oscillations have a wavenumber k of about $|\Delta M^2_{32}|$ and so the peak structure of the transform at $k \sim |\Delta M^2_{32}|$ is sensitive to these oscillations and thus the hierarchy. In Refs. [13, 15] it was shown that the heights of the peaks can be combined into two real numbers RL and PV such that $RL + PV$ is positive if and only if the hierarchy is normal. In Ref. [12] two more observables were added, one mixing information from the Fourier sine and cosine transforms and one using a nonlinear Fourier transform with the same aperiodicity as the 1-3 oscillations. A neural network finds the combination of these observables which best determines the hierarchy.

An obstruction to this analysis has been described in Ref. [16], which observed that $RL + PV$ is very sensitive to the choice of model of the reactor neutrino flux and to variations of $\Delta M^2_{\mathrm{eff}}$ smaller than the precision of its determination by MINOS [17]. To eliminate this spurious dependence, we use a weighted Fourier transform in which higher energies are weighted less heavily, providing a soft cutoff. Fig. 1 shows the average values of $RL + PV$ in simulated data using a normal and a weighted Fourier transform. The normal Fourier transform leads to the fluctuating dependence found in Ref. [16], but these fluctuations are much smaller with a weighted transform. A steeper weight would further reduce the amplitude of the fluctuations, but would also reduce the average value of $RL + PV$, weakening the hierarchy dependence.

Thus we find that the spurious dependence observed in Ref. [16] can be eliminated via a simple modification of the analysis. We will now discuss a more serious challenge which afflicts otherwise promising sites for such an experiment.

As a determination of the hierarchy requires a medium baseline, the neutrino flux arriving from each reactor will necessarily be quite low. This means both that the detector must be very large, appreciably larger than current proposals for RENO 50 [8], and also that flux from multiple reactors must be used. Multiple reactors are available, especially in places like Japan, Korea and China's Guangdong province where such experiments may



FIG. 1: Simulated average values of $RL + PV$ obtained from 100,000 neutrinos observed at a baseline of 58 km assuming the numerically interpolated reactor spectra from the 1980's [18–20]. The black dotted curve uses an unweighted Fourier transform and the red solid curve uses the weight $\exp(-0.04E^2/\text{MeV}^2)$.

be built. However multiple reactors imply multiple baselines, and so $\overline{\nu}$s will arrive at the detector with their 1-3 oscillations out of phase. Neutrinos from different reactors are not coherent with each other, wavefunctions are not added, but probabilities are added and this destroys the fine structure of the spectrum whose precise measurement is essential to a determination of the hierarchy.

We now provide the most systematic analysis of this interference effect to date. Our goal is to illustrate the effect of multiple baselines on the chance of success of the experiment and on the optimal location. We will consider two 18 GW thermal capacity idealized point reactor neutrino sources with flux given by the Gaussian form of Ref. [20], bearing in mind that a realistic flux with a weighted transform will lead to similar results. By defining an effective baseline difference, our results can be applied to configurations with many reactors. The reactor flux is normalized such that, including ν oscillations, at 58 km each 18 GW complex yields 25,000 $\overline{\nu}_e$ in 3 years.

In Fig. 2 we display the probability p of successfully determining the hierarchy with 3 and 6 years of live time for various combinations of baselines. The solid and dashed curves are analyses using respectively $RL + PV$ and a neural network optimizing 24 coefficients corresponding to the 4 hierarchy indicators of Ref. [12] and to 6 weights: $e^{-0.08E^2}, e^{-0.02E^2}, e^{-0.04(E-3.6)^2}, e^{-E/8}, e^{-0.1(E-5.25)^2}$ and $e^{-(E-3.6)^3/100}$.

The first two panels are the 6 and 3 year probabilities of success for baseline differences between 0 and



FIG. 2: Chance of success, p, after 6 years (first and third panels) and 3 years (second and fourth panels) of live time, with neutrinos from two 18 GW reactor complexes at distinct baselines. The horizontal axis is the shortest baseline, the color is the difference between baselines. On the first two panels these are 0 (black), 500 m (red), 1 km (blue), 1.5 km (purple) and 2 km (green). On the last two panels these are 5 km (black), 10 km (red), 15 km (blue) and 20 km (purple). The solid curves use $RL + PV$ and the dashed curves a neural network.

2 km. A 2 to 5 km baseline difference causes a lower chance of success. With these analysis methods, a 95% chance of a successful hierarchy determination in 6 years is only possible if the baseline difference is appreciably less than 1 km, about the effective difference for the BaiMianShi and Mudeungsan sites [21] for Daya Bay II and RENO 50. As RENO 50 uses neutrinos from only 18 GW of reactors, the times reported here must be doubled, in addition to the factor of four discussed earlier.

The last two panels illustrate that, for baseline differences of 5 to 20 km, the far detector is a background. As a result a larger baseline difference and a shorter baseline to the near reactor both increase the signal/background ratio and so *p*. These strong backgrounds reduce the optimal baseline.

Interference and flux from more distant reactors is a big problem for differences above 500 meters, arising when a detector is not perpendicular to a long linear array like the Daya Bay - Ling Ao complex. Sites such as BaiYunZhang are perpendicular and so enjoy identical baselines, but use flux from only a single array. In particular Daya Bay II sites such as BaiMianShi are not competitive. Reno 50 will face similar problems, as the largest mountains are at a 45 degree angle to the reactor array.

The minimal baseline difference at potential Daya Bay II sites that use flux from multiple reactor complexes is about 500 meters, corresponding to the area around the Dongkeng site of Ref. [21] which uses flux from the Taishan and Yangjiang complexes. Thus we find that the best case probability of determining the hierarchy is about 98%. This result depends upon the values of the neutrino mixing parameters, in general we have found that a 1σ increase in $\sin^2(\theta_{13})$ or $|\Delta M^2_{21}|$ or a 1σ decrease in $|\Delta M^2_{32}|$ can improve the hierarchy determination by $0.1\sigma - 0.3\sigma$.

The *disappearance* channel at NOνA may provide a $1-2\%$ determination of the atmospheric effective mass of Refs. [7, 22]. For $\delta \sim \pi$, this mass differs from the high energy reactor effective mass (2) by about 1.5% and from the low energy reactor effective mass $|\Delta M^2_{31}|$ by nearly 3%. As the sign of these differences depends upon the hierarchy, NOνA disappearance data can improve the hierarchy determination at Daya Bay II. However there is no such advantage if $\delta \sim 0$, as the atmospheric effective mass would be nearly equal to that of Eq. (2), which will measured more precisely at Daya Bay II.

Acknowledgement

We would like to thank C. Bonati for useful discussions and suggestions. JE is supported by the Chinese Academy of Sciences Fellowship for Young International Scientists grant number 2010Y2JA01. EC and XZ are supported in part by the NSF of China.

[1] F. P. An *et al.* [DAYA-BAY Collaboration], Phys. Rev. Lett. **108** (2012) 171803 .
[2] D. Dwyer, "Daya Bay Results," presented at Neutrino 2012.
[3] J. K. Ahn *et al.* [RENO Collaboration], Phys. Rev. Lett. **108** (2012) 191802 .
[4] R. Nichol, "Final MINOS Results," presented at Neutrino 2012.
[5] S. K. Agarwalla, S. Prakash and S. U. Sankar, arXiv:1301.2574 [hep-ph].
[6] S. T. Petcov and M. Piai, Phys. Lett. B **533** (2002) 94. S. Choubey, S. T. Petcov and M. Piai, Phys. Rev. D **68** (2003) 113006 .
[7] H. Nunokawa, S. J. Parke and R. Zukanovich Funchal, Phys. Rev. D **72** (2005) 013009 .
[8] Yeongduk Kim, "Observation of reactor neutrino disappearance at RENO," presented at νTURN 2012.
[9] Y. Wang, "Daya Bay II: The Next Generation Reactor Neutrino Experiment" presented at NuFact 2012.
[10] K. Nakamura *et al.* [Particle Data Group Collaboration], J. Phys. G **37** (2010) 075021.
[11] A. Gando *et al.* [KamLAND Collaboration], Phys. Rev. D **83** (2011) 052002.
[12] E. Ciuffoli, J. Evslin and X. Zhang, JHEP **1303** (2013) 016.
[13] L. Zhan, Y. Wang, J. Cao and L. Wen, Phys. Rev. D **78** (2008) 111103 .
[14] J. Learned, S. T. Dye, S. Pakvasa and R. C. Svoboda, Phys. Rev. D **78** (2008) 071302 .
[15] L. Zhan, Y. Wang, J. Cao and L. Wen, Phys. Rev. D **79** (2009) 073007 .
[16] X. Qian, D. A. Dwyer, R. D. McKeown, P. Vogel, W. Wang, and C. Zhang, arXiv:1208.1551 [hep-ex].
[17] R. Nichol, "Final MINOS Results," presented at Neutrino 2012.P. Adamson *et al.* [MINOS Collaboration], arXiv:1208.2915 [hep-ex].

[18] K. Schreckenbach, G. Colvin, W. Gelletly and F. Von Feilitzsch, Phys. Lett. B **160** (1985) 325.

[19] A. A. Hahn, K. Schreckenbach, G. Colvin, B. Krusche, W. Gelletly and F. Von Feilitzsch, Phys. Lett. B **218** (1989) 365.

[20] P. Vogel and J. Engel, Phys. Rev. D **39** (1989) 3378.

[21] E. Ciuffoli, J. Evslin and X. Zhang, JHEP **1212** (2012) 004.

[22] N. Okamura, Prog. Theor. Phys. **114** (2006) 1045.